

## Food and Drug Administration, HHS

## § 524.1443

no response is evident in 7 days, diagnosis and therapy should be reevaluated.

(3) For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 53 FR 12512, Apr. 15, 1988; 53 FR 27851, July 25, 1988; 64 FR 404, Jan. 5, 1999]

### § 524.1240 Levamisole.

(a) *Specifications.* The drug contains 200 milligrams of levamisole per milliliter of diethylene glycol monobutyl ether (DGME) solution.

(b) *Sponsor.* See 000061 and 053501 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.350 of this chapter.

(d) *Conditions of use. Cattle*—(1) *Amount.* 2.5 milliliters per 110 pounds (10 milligrams of levamisole per kilogram) of body weight as a single dose topically to the back of the animal.

(2) *Indications for use.* Anthelmintic effective against stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*, *Chabertia*), and lungworms (*Dictyocaulus*).

(3) *Limitations.* Conditions of constant helminth exposure may require re-treatment within 2 to 4 weeks after the first treatment. Cattle must not be slaughtered within 9 days following last treatment. Do not administer to dairy animals of breeding age. Do not treat animals before dipping or prior to exposure to heavy rain. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism, and before using in severely debilitated animals.

[52 FR 10887, Apr. 6, 1987, as amended at 53 FR 7504, Mar. 9, 1988; 62 FR 61626, Nov. 19, 1997; 67 FR 78355, Dec. 24, 2002]

### § 524.1376 2-Mercaptobenzothiazole solution.

(a) *Specifications.* The drug contains 1.3 percent 2-mercaptobenzothiazole in a suitable solvent.

(b) *Sponsor.* See 017135 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* Apply twice daily to affected area.

(2) *Indications for use.* For dogs as an aid in the treatment of hot spots

(moist dermatitis) and as first aid for scrapes and abrasions.

(3) *Limitations.* Clip hair from affected area before applying. If no improvement is seen within 1 week, consult a veterinarian.

[48 FR 15618, Apr. 12, 1983, as amended at 65 FR 50913, Aug. 22, 2000; 68 FR 33381, June 4, 2003]

### § 524.1443 Miconazole nitrate cream; miconazole nitrate lotion; miconazole nitrate spray.

(a) *Specifications.* (1) The cream contains 23 milligrams of miconazole nitrate (equivalent to 20 milligrams of miconazole base) per gram.

(2) The lotion contains 1.15 percent of miconazole nitrate (equivalent to 1 percent miconazole base).

(3) The spray product consists of a dispensing container, sprayer pump assembly, and lotion which contains 1.15 percent of miconazole nitrate (equivalent to 1-percent miconazole base).

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter for use of cream, lotion, and spray; see No. 051259 in § 510.600(c) of this chapter for use of lotion and spray.

(c) *Conditions of use.* (1) Miconazole nitrate is an antifungal agent for topical treatment of infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*.

(2) Apply once daily by rubbing into or spraying a light covering on the infected site and the immediate surrounding vicinity. Continue treatment for 2 to 4 weeks until infection is completely eradicated as determined by appropriate laboratory examination.

(3) Accurate diagnosis of infecting organism is essential. Identify by microscopic examination of a mounting of infected tissue in potassium hydroxide solution or by culture on an appropriate medium.

(4) If no improvement is observed in 2 weeks, reevaluate diagnosis and therapy.

(5) Avoid contact with eyes since irritation may result.